

Diseases Associated With Containerized Seedling Soil Mixes

R. L. James

Plant pathologist, USDA Forest Service, Northern Region, Cooperative Forestry and Pest Management, Missoula, MT

Although most peat-vermiculite mixes are relatively pathogen free, disease organisms can be introduced into containers on seed and plant debris or in irrigation water. Major root pathogens of containerized conifers include species of *Fusarium*, *Pythium*, and *Phytophthora*. *Fusarium* spp. are the most common and produce several different signs of disease, including stunting, chlorosis, and needle tip dieback, and well as killing some seedlings. Composted tree bark included in the medium will suppress many pathogenic fungi.

Production of containerized conifer seedlings by northern Rocky Mountain nurseries is increasing, and as it does so, associated disease problems have become more important. The most serious diseases of containerized seedlings are foliage and stem blights (24). Foliage pathogens spread rapidly, and environmental conditions within greenhouses are often ideal for infection and buildup of pathogens (2, 25).

Damping-off and root diseases may also be important in container operations. Most root pathogens are probably introduced either on contaminated seed or from infected plant debris within or adjacent to greenhouses (1, 9, 14, 17). In general, most container soil

mixes are relatively pathogen free (25). However, some growers have used soil mixes containing sufficient pathogen populations to cause disease.

Most soil mixes for containerized conifers contain vermiculite or perlite incorporated with sphagnum peat. This type of mix is usually well drained and acidic, two factors that help reduce diseases (11, 20). Peat-vermiculite mixes are also lightweight, uniform in composition, relatively inexpensive, and readily available; they have high water-holding capacity and their acidic nature is conducive to growing conifers (20). Soil mixes with a pH of 4.5 to 6.0 are best for proper growth of seedlings and reduced incidence of disease (16).

Diseases

Major groups of pathogens associated with root diseases of containerized seedlings are species of *Fusarium* and water molds such as *Pythium* and *Phytophthora* (18). Although water molds may be seedborne (10), they are more often introduced into container nurseries through contaminated irrigation water (17). These fungi cause disease on very young seedlings and are favored by poorly drained soil mixes and prolonged wet conditions within greenhouses.

Root diseases associated with *Fusarium* are usually more common than those associated with

Pythium or *Phytophthora*. These fungi may colonize seeds (7, 9, 19), either causing damping-off shortly after seedlings emerge or killing older seedlings. Several species of *Fusarium* are important causes of root disease of containerized conifers. These include *F. oxysporum* Schlecht. (7, 9), *F. solani* (Mart.) Sacc (7), and *F. moniliforme* Sheld. (17). These pathogens may cause chlorosis (11), stunting (22, 25), and needle tip dieback (8), as well as seedling mortality. *Fusarium* often produces spores on structures called sporodochia at the base of infected seedlings (22). These spores may spread to nearby seedlings and cause infection during watering (19). *Fusarium* may occur within peat-vermiculite mixes (8), but disease development is usually restricted if the mix is acidic (pH less than 6.0).

Disease Control

Root diseases in containerized conifer nurseries are usually sporadic, cause little damage, and do not require specific control measures. However, if disease levels are high, several procedures can help reduce losses.

Seeds should be as free of pathogens as possible. Seeds collected directly from the tree are usually less contaminated than seeds collected from the ground or squirrel caches (24). Seeds can easily be treated before sowing

to remove surface-contaminating fungi. A continuous tap water rinse for 48 hours is usually effective in removing most seed-coat fungi (7, 20). Seeds can also be treated with hydrogen peroxide or fungicides, although some effects on germination may occur (7, 24, 25).

Greenhouses should be kept clean to reduce damage from all diseases. Plant debris should be removed periodically, and benches and walls sterilized between crops (25). Diseased seedlings should be removed as soon as they are discovered (11). A noncontaminated water supply is also important (9, 17).

Soil mixes suspected of containing high populations of pathogens should be treated to reduce or eliminate these pathogens. Chemicals used to sterilize soil mixes include formaldehyde, chloropicrin, methyl bromide, and metham (Vapam) (20). The most widely used system of soil mix sterilization is heating with steam to about 82 °C (180 °F) for 30 minutes (2, 4). This will kill most harmful bacteria, fungi, nematodes, insects, and weed seeds. *Fusarium* species are killed at even lower temperatures (57 °C or 135 °F) (4). The treated soil mix should be placed in containers and handled as little as possible to reduce chances for reinfestation by pathogens (2).

Fungicides applied after root disease symptoms appear may not always be effective (24). Fun-

gicides added to soil mixes may retard seedling growth (25). If fungicides are to be used, they should be applied as a drench immediately after sowing (19). Benomyl may control *Fusarium* and ETMT (Truban) may control *Pythium*. However, because of their uncertain effectiveness, fungicides should only be used when other control measures fail.

Another approach to controlling root diseases of containerized plants is to use composted tree bark in the soil mix. Composted bark has replaced peat in soil mixes for several ornamental species grown in containers (5). One of the major advantages of composted tree bark is that it suppresses several important plant pathogens, including *Phytophthora* (6, 22, 23), *Fusarium* (21), and *Rhizoctonia* (12, 13).

Composting is a process of partially decomposing conifer or hardwood bark to produce a more absorptive, uniform material. The process includes a thermophilic phase, during which high temperatures (40 to 80 °C) kill most organisms, and a stabilization phase, during which the rate of decomposition decreases, temperatures decline, and microorganisms, some of which are antagonistic to plant pathogens, recolonize the compost (5). Most growers use a 4:1 (v/v) mixture of bark and peat as the organic component of the soil mix. This ratio results in almost

complete suppression of root diseases without the need for sterilizing the soil mix or applying fungicide.

There are three major mechanisms of root pathogen suppression from composted bark. Bark particles are generally coarser than peat, resulting in improved aeration, which is less conducive to disease occurrence (5). Composted bark supports high levels of antagonistic organisms, whereas peat does not (13). Also, water extracts from composted bark have fungicidal properties (6, 12). Composted tree bark has been used effectively in soil mixes to control plant pathogens in several ornamental plant industries. This approach should also be considered in containerized conifer seedling operations.

Literature Cited

1. Armson, K.A.; Sadrieka, V. Forest tree nursery soil management and related practices. Toronto: Ontario Ministry of Natural Resources; 1974. 143 p.
2. Baker, K.F.; Olsen, S.M. Soil steaming. California State Florist Association Bulletin 8(9): 1-10; 1959.
3. Chef, D.G. Suppression of *Fusarium* wilt of chrysanthemum in composted hardwood bark. M.S. Thesis. Columbus: Ohio State University; 1977. 38 p.
4. Hartmann, H.T.; Kester, D.E. Plant propagation. 3d ed. Englewood Cliffs, NJ: Prentice-Hall; 1959. 648 p.
5. Hoitink, H.A.J. Composted bark, a lightweight growth medium with fungicidal properties. Plant Disease 64: 142-147; 1980.

6. Hoitink, H.A.J.; Van Doren, D.M., Jr.; Schmitthenner, A.F. Suppression of *Phytophthora cinnamomi* in composted hardwood bark potting medium. *Phytopathology* 67: 561-565; 1977.
7. James, R.L. Fusarium root disease of containerized seedlings at the Montana State Nursery, Missoula. Missoula, MT: U.S. Department of Agriculture, Forest Service, Northern Region; 1983. 9 p.
8. James, R.L. Needle tip dieback of containerized Douglas-fir seedlings at the Coeur d' Alene Nursery, Idaho. Missoula, MT: U.S. Department of Agriculture, Forest Service, Northern Region; 1983; 9 p.
9. James, R.L. Occurrence of *Fusarium* on Douglas-fir seed from the Coeur d' Alene Nursery. Missoula, MT: U.S. Department of Agriculture, Forest Service, Northern Region; 1983. 11 p.
10. James, R.L.; Genz, D. Evaluation of fungal populations on ponderosa pine seed. Rep. 82-22. Missoula, MT: U.S. Department of Agriculture, Forest Service, Northern Region; 1982. 21 p.
11. Landis, T.D. Fusarium root disease of containerized tree seedlings—Colorado State Forest Service Nursery. Biol. Eval. R2-76-16. Lakewood, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Region; 1976. 6 p.
12. Nelson, E.B.; Hoitink, H.A.J. Factors affecting suppression of *Rhizoctonia solani* in container medium. *Phytopathology* 72: 275-279; 1982.
13. Nelson, E.B.; Hoitink, H.A.J. The role of microorganisms in the suppression of *Rhizoctonia solani* in container media amended with composted hardwood bark. *Phytopathology* 73: 274-278; 1983.
14. Pawuk, W.H. Damping-off of container-grown longleaf pine seedlings by seedborne fusaria. *Plant Disease Reporter* 62: 82-84; 1978.
15. Pawuk, W.H. Fungicide coverings affect the germination of southern pine seeds. *Tree Planters' Notes* 30(1): 3-4; 1979.
16. Pawuk, W.H. Potting media affect growth and disease development of container-grown southern pines. Res. Note SO-268. Atlanta, GA: U.S. Department of Agriculture, Forest Service; 1981 4 p.
17. Pawuk, W.H. Diseases of container-grown southern pine seedlings and their control. In: Guldin, R.W.; Barnett, J.P. (eds.). *Proceedings, Southern Containerized Forest Tree Seedling Conference*; 1981; Savannah, GA. Gen. Tech. Rep. SO-37. Atlanta: U.S. Department of Agriculture, Forest Service; 1982: 47-50.
18. Pawuk, W.H.; Barnett, J.P. Root rot and damping-off of container-grown southern pine seedlings. In: Tinus, R.W.; Stein, W.I.; Balmer, W.E. (eds.). *Proceedings, North American containerized forest tree seedling symposium*; 1974; Denver, CO: Great Plains Agricultural Council Publ. 68. Washington, DC: U.S. Government Printing Office; 1974: 173-176.
19. Peterson, G.W. Disease problems in the production of containerized forest tree seedlings in North America. In: Tinus, R.W.; Stein, W.I.; Balmer, W.E. (eds.). *Proceedings, North American containerized forest tree seedling symposium*; 1974; Denver, CO: Great Plains Agricultural Council Publ. 68. Washington, DC: U.S. Government Printing Office; 1974: 170-172.
20. Phipps, H.M. Growing media affect size of container-grown red pine. Res. Note NC-165. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1974. 4 p.
21. Sekiguchi, A. Control of *Fusarium* wilt on Chinese yam. Annu. Rep., Department of Plant Pathology and Entomology, Nagano Vegetable and Floriculture Experiment Station, Nagano, Japan. 1: 10-11, 1977.
22. Spencer, S.; Benson, D.M. Root rot of *Aucuba japonica* caused by *Phytophthora cinnamomi* and *P. citricola* and suppressed with bark media. *Plant Disease* 65: 918-921; 1981.
23. Spencer, S.; Benson, D.M. Pine bark, hardwood bark compost, and peat amendment effects on development of *Phytophthora* spp. and lupine root rot. *Phytopathology* 72: 346-351, 1982.
24. Sutherland, J.R.; Lock, W.; Benson, L.E. Diseases and insects and their management in container nurseries. In: Scarratt, R., Jr.; Glerum, C.; Phexman, C.A. (eds.). *Proceedings, Canadian containerized tree seedling symposium*; 1981; Toronto. Canadian-Ontario Joint Forestry Research Committee Symposium Proceedings O-P-10. Sault Ste. Marie, ON: Canadian Forest Service, Department of the Environment, Great Lakes Forest Research Center; 1982: 215-223.
25. Tinus, R.W.; McDonald, S.E. How to grow tree seedlings in containers in greenhouses. Gen. Tech. Rep. RM-60. Lakewood, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station; 1979. 256 p.