

# 1972 Author Index

(Tree Planters' Notes, Vol. 23, 1-4)

**Note:** First figure is issue number; second is page.

- Alford, Lee T., and DeBell, Dean S.**  
Sprouting Characteristics and Cutting Practices Evaluated for Cottonwoods—Could Lead to Higher Yields 4:1
- Barnett, James P.**  
Drying and Sorting Stratified Loblolly Pine Seeds Reduces Dormancy 3:10
- Campbell, Robert K., and Wilson, Boyd C.**  
Seedbed Density Influences Height, Diameter, and Dry Weight of 3-0 Douglas Fir 2:1
- Campbell, W. A., Hendrix, F. F., and Powell, W. M.**  
Pythium and Nematode Species Implicated in Root Rot 1:5
- Chester, G., and Wilde, S. A.**  
Air Humidity, Ground Water Extend Growing in Wisconsin 3:19
- Day, M. W., White, D. P., and Wright, J. W.**  
Fertilizer Application Can Improve Red Pine Seed Production 1:25
- DeBell, Deans S., and Alford, Lee T.**  
Sprouting Characteristics and Cutting Practices Evaluated for Cottonwoods—Could Lead to Higher Yields 4:1
- Dickson, Richard, and McBride, Joe R.**  
Giberellic, Citric Acids and Stratification Enhance White Ash Germination . . . Growth Inhibitor Reversed 3:1
- Dorn, Donald E., and Pitcher, John A.**  
Geographic Source Differences Noted in Black Cherry Seed Weight, Germination 3:7
- Edgren, James W.**  
Field Survival Poor for Stored and Heeled-In Ponderosa Pine Seedlings 1:19
- Ellis, Gilbert R.**  
Plastic Mesh Tubes Constrict Black Walnut Root Development After Two Years 3:27
- Forbes, Donovan C.**  
Black Cherry Seed Germination: A Comparison of Seeds Collected at Different Stages of Maturity 3:5
- Gilbert, R. C. A.**  
Controlling Competing Vegetation . . . Sprayer-Equipped Planter Applies Herbicides at Planting Time 2:8
- Goddard, R. E., Hickman, J. G., and Smith, W. H.**  
Techniques for Improved Grafting 4:13
- Hallman, R. G., and Stoleson, R. M.**  
Helicopter Seeders Can Damage Conifer Seed 3:29
- Hatchell, G. E., and Langdon, O. Gordon**  
Carolina Studies Offer Possibilities of Increasing Pine Growth on Savannah Sites 2:5
- Hendrix, F. F., Powell, W. M., and Campbell, W. A.**  
Pythium and Nematode Species Implicated in Root Rot 1:5
- Hickman, J. G., Goddard, R. E., and Smith, W. H.**  
Techniques for Improved Grafting 4:13
- Hocking, Drake**  
Nursery Practices in Cold Storage of Conifer Seedlings in Canada and the United States: A Survey 2:26
- Hocking, D., and Ward, B.**  
Late Lifting and Freezing in Plastic Bags Improve White Spruce Survival After Storage 3:24
- Hook, Donal D., Kormanik, Paul, P., and McAlpine, R. G.**  
Horizontal Planting of Sycamore Cuttings: Better and Cheaper Method of Establishing High-Density Stand Sought 2:5
- Hooper, Ralph M., and Olson, David F., Jr.**  
Northern Red Oak Plantings Survive Well in Southern Appalachians, But Growth is Slow 1:16
- Huntzinger, Harold J.**  
Direct Seeding Black Cherry: Some Recommendations for the Allegheny Plateau 1:10
- Irish, J. Jack, and Nyland, Ralph D.**  
Unusual Ice Damage Suggests Extra Care Needed in Overwinter Cold Storage . . . "Windowpaning" Injures Branch Tips 3:13
- Jackson, H.**  
British Nursery Reports Success with Finnish Transplanting Technique 4:26
- Kormanik, Paul P., McAlpine, Robert G., and Hook, D. D.**  
Horizontal Planting of Sycamore Cuttings: Better and Cheaper Method of Establishing High-Density Stand Sought 2:5
- Langdon, O. Gordon, and Hatchell, G. E.**  
Carolina Studies Offer Possibilities of Increasing Pine Growth on Savannah Sites 2:19

|                                                                                                                                                          |             |                                                                                                                        |             |                                                                                                                                                          |             |
|----------------------------------------------------------------------------------------------------------------------------------------------------------|-------------|------------------------------------------------------------------------------------------------------------------------|-------------|----------------------------------------------------------------------------------------------------------------------------------------------------------|-------------|
| <b>Lock, W., Sluggett, L. J., and Sutherland, Jack R.</b>                                                                                                |             | <b>Oak and Loblolly Pine</b>                                                                                           | <b>1:20</b> | <b>Techniques for Improved Grafting</b>                                                                                                                  | <b>4:13</b> |
| <b>Corky Root Disease Observed on Two Spruce Species and Western Hemlock</b>                                                                             | <b>4:18</b> | <b>Peterson, Glenn W.</b>                                                                                              |             | <b>Stein, William I., and Ouston, Peyton W.</b>                                                                                                          |             |
| <b>Losche, Craig K.</b>                                                                                                                                  |             | <b>Chemical Control of Phomopsis Blight of Junipers: A Search for New Methods</b>                                      | <b>3:3</b>  | <b>Coating Materials Protect Douglas-Fir, Noble Fir Seedlings Against Drying Conditions . . . Not Recommended for Extended Storage—More Tests Needed</b> | <b>3:21</b> |
| <b>Vertical Tillage Fails to Improve Black Walnut Seedling Growth on Poorly Drained Site</b>                                                             | <b>1:7</b>  | <b>Phares, Robert E., and White, Gordon</b>                                                                            |             | <b>Stoleson, R. M., and Hallman, R. G.</b>                                                                                                               |             |
| <b>Lyle, E. S., Jr.</b>                                                                                                                                  |             | <b>Large Stock, Deep Planting Improve Cottonwood Growth in Mississippi</b>                                             | <b>4:16</b> | <b>Helicopter Seeders Can Damage Conifer Seed</b>                                                                                                        | <b>3:29</b> |
| <b>Diagnosing Mineral Deficiency by Foliar Fertilization</b>                                                                                             | <b>1:23</b> | <b>Pitcher, John A.</b>                                                                                                |             | <b>Sutherland, Jack R., Sluggett, L. J., and Lock, W.</b>                                                                                                |             |
| <b>McAlpine, Robert G., Hook, D. D., and Kormanik, P. P.</b>                                                                                             |             | <b>Repeated Fertilization Fails to Stimulate Cone Production in Young Loblolly Pine Stand in Illinois</b>              | <b>4:28</b> | <b>Corky Root Disease Observed on Two Spruce Species and Western Hemlock</b>                                                                             | <b>4:18</b> |
| <b>Horizontal Planting of Sycamore Cuttings: Better and Cheaper Method of Establishing High-Density Stand Sought</b>                                     | <b>2:5</b>  | <b>Pitcher, John A., and Dorn, Donald E.</b>                                                                           |             | <b>Tinus, Richard W.</b>                                                                                                                                 |             |
| <b>McBride, Joe R., and Dickson, Richard</b>                                                                                                             |             | <b>Geographic Source Differences Noted in Black Cherry Seed Weight, Germination</b>                                    | <b>3:7</b>  | <b>CO<sub>2</sub> Enriched Atmosphere Speeds Growth of Ponderosa Pine and Blue Spruce</b>                                                                | <b>1:12</b> |
| <b>Giberellic, Citric Acids and Stratification Enhance White Ash Germination: Growth Inhibitor Reversed</b>                                              | <b>3:1</b>  | <b>Plass, William T.</b>                                                                                               |             | <b>Von Reider, Michael, and Schwalm, John M.</b>                                                                                                         |             |
| <b>Mohlberg, John M.</b>                                                                                                                                 |             | <b>Fertilization Treatments Increase Black Locust Growth on Extremely Acid Surface-Mine Spoils</b>                     | <b>4:10</b> | <b>Wire-rolling Machine Saves Man-Hours</b>                                                                                                              | <b>3:28</b> |
| <b>Quick, Inexpensive Broadcast Seeding</b>                                                                                                              | <b>4:20</b> | <b>Powell, W. M., Campbell, W. A., and Hendrix, F. F., Jr.</b>                                                         |             | <b>Walters, Gerald A.</b>                                                                                                                                |             |
| <b>Nyland, Ralph D., and Irish, H. Jack</b>                                                                                                              |             | <b>Pythium and Nematode Species Implicated in Root Rot</b>                                                             | <b>1:5</b>  | <b>Pesticide Treatments on Saligna Eucalyptus, Australian Toon Seedlings Affect Dieback but Not Survival</b>                                             | <b>3:16</b> |
| <b>Unusual Ice Damage Suggests Extra Care Needed in Overwinter Cold Storage . . . "Windowpaning" Injures Branch tips</b>                                 | <b>3:13</b> | <b>Scarratt, J. B.</b>                                                                                                 |             | <b>Walters, Gerald A.</b>                                                                                                                                |             |
| <b>Olson, David F., Jr.</b>                                                                                                                              |             | <b>Container Size Affects Dimensions of White Spruce, Jack Pine Planting Stock . . . Larger Tubes Offer Advantages</b> | <b>4:21</b> | <b>Chemical Treatment of Bare-Root Saligna Eucalyptus Seedlings Offers No Advantages</b>                                                                 | <b>4:4</b>  |
| <b>Northern Red Oak Plantings Survive Well in Southern Appalachians—But Growth is Slow</b>                                                               | <b>1:16</b> | <b>Schneider, G., and White, Donald P.</b>                                                                             |             | <b>Walters, Gerald A.</b>                                                                                                                                |             |
| <b>Ouston, Peyton W., and Stein, William I.</b>                                                                                                          |             | <b>Soilless Container System Developed for Growing Conifer Seedlings</b>                                               | <b>1:1</b>  | <b>Packing Methods Studied for Australian Toon and Slash Pine Plantings</b>                                                                              | <b>4:7</b>  |
| <b>Coating Materials Protect Douglas-Fir, Noble Fir Seedlings Against Drying Conditions . . . Not Recommended for Extended Storage—More Tests Needed</b> | <b>3:21</b> | <b>Schwalm, John M., and von Reider, Michael</b>                                                                       |             | <b>Ward, B. and Hocking, D.</b>                                                                                                                          |             |
| <b>Parris, G. K.</b>                                                                                                                                     |             | <b>Wire-Rolling Machine Saves Man-Hours</b>                                                                            | <b>3:28</b> | <b>Late Lifting and Freezing in Plastic Bags Improve White Spruce Survival After Storage</b>                                                             | <b>3:24</b> |
| <b>Improperly Applied Weedkillers Can Damage or Kill</b>                                                                                                 |             | <b>Sluggett, L. J., Lock W., and Sutherland, Jack R.</b>                                                               |             | <b>White, D. P., Day, M. W., and Wright, J. W.</b>                                                                                                       |             |
|                                                                                                                                                          |             | <b>Corky Root Disease Observed on Two Spruce Species and Western Hemlock</b>                                           | <b>4:18</b> | <b>Fertilizer Application Can Improve Red Pine Seed Production</b>                                                                                       | <b>1:25</b> |
|                                                                                                                                                          |             | <b>Smith, W. H., Goddard, R. E., and Hickman, J. G.</b>                                                                |             | <b>White, Donald P., and Schneider, G.</b>                                                                                                               |             |

|                                                                     |      |                                                                                                                |      |                                                                   |
|---------------------------------------------------------------------|------|----------------------------------------------------------------------------------------------------------------|------|-------------------------------------------------------------------|
| Soilless Container System Developed for Growing Conifer Seedlings   | 1:1  | of Black Walnut Seedlings                                                                                      | 2:22 | Woessner, Ronald A.                                               |
| White, Gordon, and Phares, Robert E.                                |      | Williston, Hamlin L.                                                                                           |      | Four Hardwood Species Differ in Tolerance to Pruning              |
| Large Stock, Deep Planting Improve Cottonwood Growth in Mississippi | 4:16 | The Question of Adequate Stocking—Pine Planting Project Must Allow for Seedling Survival and Plantation Losses | 1:9  | 1:28                                                              |
| Wilde, S. A.                                                        |      | Wilson, Boyd C., and Campbell, Robert K.                                                                       |      | Woessner, Ronald A.                                               |
| Air Humidity, Ground Water Extend Growing Season in Wisconsin       | 3:19 | Seedbed Density Influences Height, Diameter and Dry Weight of 3-0 Douglas-Fir                                  | 2:1  | Weed Control by Herbicides Promotes Growth in Cottonwood Cuttings |
| Williams, Robert D.                                                 |      | Woessner, Ronald A.                                                                                            |      | 2:17                                                              |
| Root Fibrosity Proves Insignificant in Survival, Growth             |      | Epicormic Sprouting of Eastern Cottonwood Clones Affected by Pruning Date                                      | 3:12 | Wright, J. W., Day, M. W., and White, D. P.                       |
|                                                                     |      |                                                                                                                |      | Fertilizer Application Can Improve Red Pine Seed Production       |
|                                                                     |      |                                                                                                                |      | 1:25                                                              |

# 1972 Subject and Title Index

(Tree Planters' Notes, Vol. 23, 1-4)

**Note:** Listing followed by ( . . . ) indicates title.

**AIR HUMIDITY . . .**, ground water extend growing season in Wisconsin 3:19

**ALLEGHENY PLATEAU**  
Some recommendations for, — Direct seeding black cherry 1:10

**APPALACHIANS, SOUTHERN**  
Nothorn red oak plantings survive well in Southern Appalachians but growth is slow 1:16

**AUSTRALIAN TOON**  
Pesticide treatment on saligna eucalyptus, australian toon seedlings affect dieback but not survival 3:16

Australian toon and slash pine plantings,  
Packing methods studied for 4:7

**BLACK CHERRY**  
Direct seeding: some recommendations for the Allegheny Plateau 1:10

Black cherry seed germination: a comparison of seeds collected at different stages of maturity 3:5

Geographic source differences noted in black cherry seed weight 3:7

**BLACK LOCUST**  
Fertilization treatments increase growth on extremely acid surface mine spoils 4:10

**BLACK WALNUT**  
Seedlings—Root fibrosity proves insignificant in survival, growth of 2:22

Seedling growth on poorly drained site, Vertical tillage fails to improve 1:7

Root development, after two years, Plash mesh tubes constrict 3:27

**BLUE SPRUCE**  
And ponderosa pine seedlings, CO<sub>2</sub> enriched atmosphere speeds growth of 1:12

**BRITISH . . .** nursery reports success with Finnish transplanting techniques 4:26

<sup>1</sup> Subject followed by . . . indicates title listing.

**BROADCAST SEEDING**  
Quick, inexpensive 4:20

## CANADA

And the U.S., Nursery practices in cold storage of conifer seedlings in—A survey 2:26

**CAROLINA . . .** studies offer possibilities of increasing pine growth on savannah sites 2:19

## CHEMICAL

Chemical control of phomopsis blight of junipers: a search for new methods 3:3

Chemical treatment of bare-root saligna eucalyptus seedlings offers no advantages 4:4

## CITRIC ACID

Giberellic, citric acids, and stratification enhance white ash germination — Growth inhibitor reversed 3:1

## CLONES

Epicormic sprouting of eastern cottonwood, affected by pruning date 3:12

CO<sub>2</sub> . . . enriched atmosphere speeds growth of ponderosa pine and blue spruce seedlings 1:12

## COATING MATERIALS

Coating materials, protect doug-

- las-fir, noble fir seedling against drying conditions — Not recommended for extended storage — more tests needed 3:21
- COLD STORAGE**  
Of conifer seedlings, nursery practices in Canada and the United States: A survey 2:26  
“Windowpaning” injures branch tips  
Unusual ice damage suggests extra care needed in overwinter cold storage 3:13
- CONES**  
Cone production in young loblolly pine stand in Illinois, repeated fertilization fails to stimulate 4:28
- CONIFER SEED**  
Helicopter seeders can damage 3:29  
Conifer seedlings in Canada and the United States,  
Nursery practices in cold storage of: A survey 2:26  
Soiless container system developed for growing conifer seedlings 1:1
- CONTAINER**  
Container size affects dimensions of white spruce, jack pine planting stock—Larger tubes offer advantages 4:21  
Container system, Soiless, developed for growing conifer seedlings 1:1
- COTTONWOODS**  
Cottonwood clones, epicormic sprouting of eastern, affected by pruning date 3:12  
Sprouting characteristics and cutting practices evaluated for — could lead to higher yields 4:1  
Cuttings, Weed control by herbicides promotes growth of 2:17  
Cottonwood growth in Mississippi, Large stock, deep planting improve 4:16
- CORKY ROOT DISEASE . . .** observed on two spruce species and western hemlock 4:18
- CUTTINGS**  
Cottonwood, Weed control by herbicides at planting time promotes growth of 2:17  
Cutting practices evaluated for cottonwoods — Could lead to higher yields 4:1
- DIAGNOSING . . .** mineral deficiency by foliar fertilization 1:23
- DIAMETER**  
Height, and dry weight of 3-0 Douglas-fir, Seedbed density influences 2:1
- DIRECT SEEDING . . .** black cherry — Some recommendations for the Allegheny Plateau 1:10
- DISEASE**  
Corky root, observed on two spruce species and western hemlock 4:18
- DORMANCY**  
Drying and storing stratified loblolly pine seeds reinduces 3:10
- DOUGLAS FIR**  
Douglas fir, noble fir seedlings, Coating materials protect against drying conditions — Not recommended for extended storage, more tests needed 3:21  
Douglas-fir, height, diameter, and dry weight of 3-0, Seedbed density influences 2:1
- DRYING . . .** and storing stratified loblolly pine seeds reinduces dormancy 3:10
- EASTERN COTTONWOOD**  
Clones, epicormic sprouting of, affected by pruning date 3:12
- EPICORMIC . . .** sprouting of eastern cottonwood clones affected by pruning date 3:12
- FERTILIZATION**  
Repeated, fails to stimulate cone production in young loblolly pine stand in Illinois, Repeated 4:28  
Fertilization treatments increase black locust growth on extremely acid surface-mine spoils 4:10
- FIELD SURVIVAL . . .** poor for stored and heeled-in ponderosa pine seedlings 1:19
- FINLAND**  
Finnish transplanting technique, British nursery reports success with 2:26
- FOLIAR FERTILIZATION**  
Diagnosing mineral deficiency by 1:23
- FOUR HARDWOOD SPECIES . . .** differ in tolerance to pruning 1:28
- FREEZING**  
In plastic bags, Late lifting and, improve white spruce survival after storage 3:24
- GEOGRAPHIC . . .** source differences noted in black cherry seed weight, germination 3:7
- GERMINATION**  
Black cherry seed, Comparison of seeds collected at different stages of maturity 3:5  
Germination, black cherry seed weight and, Geographic source differences noted in 3:7
- GIBERELLIC, . . .** citric acids and stratification enhance white ash germination — Growth inhibitor reversed 3:1
- GROUND WATER**  
Air humidity and, extend growing season in Wisconsin 3:19
- GROWTH**  
Growth inhibitor reversed, Gibberellic, citric acids and stratification enhance white ash germination 3:1  
Growth, cottonwood, in Mississippi, Large stock, deep planting improve 4:16  
Growth is slow, Northern red oak plantings survive well in southern Appalachians but 1:16  
Growth, survival of black walnut seedlings, Root fibrosis proves insignificant in 2:22  
Growth of cottonwood cuttings, Weed control by herbicides promotes 2:17  
Growth of ponderosa pine and

- blue spruce seedlings, CO<sub>2</sub> enriched atmosphere speeds 1:12  
 Black locust, Fertilization treatments increase, on extremely acid surface-mine spoils 4:10  
 Black walnut seedling, Vertical tillage fails to improve on poorly drained site 4:10
- HARDWOODS**  
 Four hardwood species differ in tolerance to pruning 1:28
- HEELED-IN**  
 Heeled-in ponderosa pine seedlings,  
 Field survival poor for stored and 1:19
- HEIGHT**  
 Height, diameter, and dry weight of 3-0 Douglas-fir, Seedbed density influences 2:1
- HELICOPTERS**  
 Helicopter seeders can damage conifer seed 3:29
- HERBICIDES**  
 Weed control by, Promotes growth of cottonwood cuttings 2:17
- HORIZONTAL PLANTING . . .**  
 of sycamore cuttings:  
 Better and cheaper method of establishing high-density stand sought 2:5
- ICE DAMAGE . . .** suggests extra care needed in overwinter cold storage — "Windowpaning" injures branch tips 3:13
- ILLINOIS**  
 Repeated fertilization fails to improve cone production in young loblolly pine stand in 4:28
- IMPROPERLY APPLIED WEED KILLERS . . .** can damage or kill oak and loblolly pine 12:20
- JACK PINE**  
 And white spruce planting stock, Container size affects dimensions of—larger tubes offer advantages 4:21
- JUNIPERS**  
 Chemical control of phomopsis blight of: A search for new methods 3:3
- LARGE STOCK . . .**, deep planting improve cottonwood growth in Mississippi 4:16
- LOBLOLLY PINE**  
 Oak, and, Improperly applied weed killers can damage or kill 1:20  
 Loblolly pine seed, drying and storing stratified, reinduces dormancy 3:10  
 Loblolly pine stand in Illinois, Repeated fertilization fails to stimulate cone production in young 4:28
- MACHINE**  
 Wire-rolling, saves man-hours 3:28
- MINERAL**  
 Mineral deficiency, Diagnosing by foliar fertilization 1:23
- MISSISSIPPI**  
 Large stock, deep planting improve cottonwood growth in 4:16
- NEMATODE . . .** and pythium species implicated in root rot 1:5
- NOBLE FIR**  
 Douglas fir and noble fir seedlings, Coating materials protect against drying conditions — not recommended for extended storage — more tests needed 3:21
- NORTHERN RED OAK**  
 Plantings survive well in southern Appalachians—but growth is slow 1:16
- NURSERY**  
 British, reports success with Finnish transplanting technique 4:26  
 . . . practices in cold storage of conifer seedlings in Canada and the United States: A survey 2:26
- OAK**  
 Oak and loblolly pine, Improperly applied weedkillers can damage or kill 1:20  
 Oak, northern red, Plantings survive well in Southern Appalachians—but growth is slow 1:16
- PACKING . . .** methods studied for Australian toon and slash pine plantings 4:7
- PESTICIDE**  
 Pesticide treatments on saligna eucalyptus, australian toon seedlings affect dieback but not survival 3:16
- PINE**  
 Pine growth of savannah sites, Carolina studies offer possibilities of increasing 2:19  
 Loblolly, Improperly applied weedkillers can damage or kill oak and 1:20  
 . . . planting project must allow for seedling survival and plantation losses—the question of adequate stocking 1:9  
 Pine, red, Fertilizer application can improve seed production of 1:25
- PLANTING**  
 Horizontal, of sycamore cuttings —better and cheaper method of establishing high-density stand sought 2:5  
 Plantings, Packing methods studied for australian toon and slash pine 4:7  
 Planting stock — Container size affect dimensions of white spruce, jack pine, larger tubes offer advantages 2:21  
 Northern Red Oak, survive well in Southern Appalachians — but growth is slow 1:16
- PLASTIC**  
 . . . mesh tubes constrict black walnut root development after 2 years 3:27  
 Bags, late lifting and freezing in, improve white spruce survival after storage 3:24
- PONDEROSA PINE**  
 Ponderosa pine and blue spruce seedlings, CO<sub>2</sub> enriched atmosphere speeds growth of 1:12  
 Ponderosa pine seedlings, Field survival poor for stored and heeled-in 1:19
- PRUNING**  
 Date, Epicormic sprouting of

- eastern cottonwood clones affected by 3:12  
Pruning, Four hardwood species differ in tolerance to 1:28
- PYTHIUM** . . . and nematode species implicated in root rot 1:5
- QUICK** . . . , inexpensive broadcast seeding 4:20
- RED PINE**  
Red pine seed production, fertilizer application can improve 1:25
- ROOT**  
Root rot, Pythium and nematode species implicated in 1:5  
. . . fibrosity proves insignificant in survival, growth of black walnut seedlings 2:22
- SALIGNA EUCALYPTUS**  
Saligna eucalyptus, Australian toon seedlings, Pesticide treatments on, affect dieback but not survival 3:16  
Saligna eucalyptus seedlings, no advantages, Chemical treatment of bare-root 4:4
- SAVANNAH**  
Savannah sites, Carolina studies offer possibilities of increasing pine growth on 2:19
- SEEDBED** . . . density influences height, diameter, and dry weight of 3-0 Douglas-fir 2:1
- SEED**  
Production, red pine, Fertilizer application can improve 1:25  
Weight, germination, black cherry Geographic source differences noted in 3:7
- SEEDING**  
Direct, black cherry: Some recommendations for the Allegheny Plateau 1:10
- SEEDLING**  
Seedling, black walnut, Vertical tillage fails to improve growth of, on poorly drained site 1:7  
Seedlings, bare-root saligna eucalyptus, Chemical treatment of, offers no advantages 4:4  
Seedlings, black walnut, survival and growth of, Root fibrosity proves insignificant in 2:22  
Seedling survival and plantation losses, pine planting project must allow for—the question of adequate stock 1:9
- SLASH PINE**  
Slash pine plantings, Packing methods studied for Australian toon and 4:7
- SOILLESS** . . . container system developed for growing conifer seedlings 1:1
- SPRAYER**  
. . . -equipped planter applies herbicides at planting time—controlling competing vegetation 2:8
- SPROUTING**  
. . . characteristics and cutting practices evaluated for cottonwoods—Could lead to higher yields 4:1
- SPRUCE**  
Spruce species, two, and western hemlock, Corky root disease observed on 4:18
- STOCKING**  
Stocking, The question of adequate — Pine planting projects must allow for seedlings survival and plantation losses 1:9
- STORAGE**  
Cold, of conifer seedlings, Nursery practices in Canada and U.S.: A survey 2:26  
Overwinter, Unusual ice damage suggests extra care needed in—“Windowpaning” injures branch tips 3:13
- STRATIFICATION**  
Stratification, Giberellic, citric acids and, enhance white ash germination — Growth inhibitor reversed 3:1
- SURFACE-MINE**  
Spoils, Fertilization treatments increase black locust growth on extremely acid 4:10
- SURVIVAL**  
After storage, white spruce, Late lifting and freezing in plastic bags improve 3:24
- Survival, growth of black walnut seedlings, Root fibrosity proves insignificant in 2:22
- SYCAMORE**  
Horizontal planting of — could lead to higher yields 4:1
- TILLAGE**  
Vertical, fails to improve black walnut seedling growth on poorly drained site 1:7
- TRANSPLANTING**  
British nursery reports success with Finnish technique 4:26
- TUBES**  
Plastic mesh, constrict black walnut root development after 2 years 3:27
- VERTICAL** . . . tillage fails to improve black walnut seedling growth on poorly drained sites 1:7
- WEED**  
. . . control by herbicides promotes growth of cottonwood cuttings 2:17
- WEEDKILLERS**  
Improperly applied, can damage or kill oak and loblolly pine 12:20
- WESTERN HEMLOCK**  
Western hemlock, Corky root disease observed on two spruce species and 4:18
- WHITE ASH**  
White ash germination—Giberellic, citric acids and stratification enhance — Growth inhibitor reversed 3:1
- WHITE SPRUCE**  
White spruce, jack pine, planting stock, Container size affects dimensions of—Larger tubes offer advantages 4:21  
White spruce survival after storage, Late lifting and freezing in plastic bags improve 3:24
- WIRE-ROLLING** . . . machine saves man-hours 3:28
- WISCONSIN**  
Air humidity, ground water extend growing season in 3:19

