

Tree Planters' Notes Index for Volume 38 (1987)

Subject Index

Abies

fraseri. See Fraser fir

procera. See Noble fir

Alnus rubra. See Red alder

Animal damage/control. See also Insect damage/
control

deer in Douglas-fir plantations, repellants
for 38(3):22-25, Summer 1987

elk in Douglas-fir plantations, repellants for
38(3):22-25, Summer 1987

nutria and baldcypress seedlings 38(3):26-29,
Summer 1987

pocket gopher and black cherry, northern red oak
38(4):31-36, Fall 1987

Antitranspirant

Cloud Cover 38(4):23-26, Fall 1987

Vapor Gard 38(1):26-30, Winter 1987

Wilt-pruf NCF 38(4):23-26, Fall 1987

X2-1337 38(4):23-26, Fall 1987

Austrian pine

bleach surface-sterilized seeds and Fusarium and
damping-off 38(3):18-21, Summer 1987

seed germination 38(3):18-21, Summer 1987

Autumn olive

germination, stratification/temperature

recommendations 38(1):14-17, Winter 1987

seed treatment 38(1):14-17, Winter 1987

Baldcypress

nutria control trial 38(3):26-29, Summer 1987

Vexar plastic seedling protector 38(3):26-29, Summer
1987

Benomyl

loblolly pine seedlings, cold-stored, effect on
38(4):19-22, Fall 1987

sycamore seedlings, cold-stored, effect on
38(1):26-30, Winter 1987

Biomass production

aboveground, of black cottonwood, red alder,
Eucalyptus 38(4):27-30, Fall 1987

Birch Quik-Tach Scalper

advantages and disadvantages 38(1):3-5, Winter 1987

Black cherry

adaptability to northern California and 15-year

survival/growth results 38(4):31-36, Fall 1987

competition with grasses, forbs 38(4):31-36, Fall 1987

pocket gopher, deer, sawfly damage 38(4):3136, Fall
1987

Black cottonwood

biomass production, aboveground 38(4):27-30, Fall
1987

cutting survival 38(4):27-30, Fall 1987

early height growth 38(4):27-30, Fall 1987

frost tolerance 38(4):27-30, Fall 1987

Black spruce

antitranspirants, film-forming, effect on seedlings
38(4):23-26, Fall 1987

plant moisture stress of seedlings 38(4):23-26, Fall 1987

seedlings, outplanted, survival 38(4):23-26, Fall 1987

Black walnut

adaptability to northern California and 15-year

survival/growth results 38(4):31-36, Fall 1987

soil moisture 38(4):31-36, Fall 1987

Bleach, 40% laundry

surface sterilization of seeds 38(3):18-21, Summer
1987

Brown spot

benomyl 38(4):19-22, Fall 1987

longleaf pine, effect of seedling size on
38(3):16-17, Summer 1987

Bud dormancy. See Dormancy, bud

Carya illinoensis. See Pecan

Chemicals. See specific names; Fungicides

Chilling. See Seedling storage, cold/frozen

Christmas trees. See Fraser fir

Clearwater Yarder

test results and recommendations 38(1):6-10, Winter
1987

Cloud Cover

on black spruce seedlings 38(4):23-26, Fall 1987

Cold hardiness

cold-stored Douglas-fir, lodgepole pine, interior spruce seedlings 38(2):11-15, Spring 1987

Eucalyptus macarthurii and *E. viminalis* in Pacific Northwest 38(4):27-30, Fall 1987

method, simple, for evaluating whole plant 38(2):16-18, Spring 1987

Cold storage. See Seedling storage, cold/frozen

Compost. See also Soil treatment

sawdust-sludge, effect on 3 conifer species 38(2):22-27, Spring 1987

Container-grown seedlings

black spruce 38(4):23-26, Fall 1987

Douglas-fir 38(1):18-21, Winter 1987

Engelmann spruce 38(1):18-21, Winter 1987

Fraser fir 38(2):19-21, Spring 1987

limber pine 38(1):18-21, Winter 1987

lodgepole pine 38(1):18-21, Winter 1987

ponderosa pine 38(1):18-21, Winter 1987

tomato 38(1):24-25, Winter 1987

Cost. See Economics

Cuttings. See also Seedling growth/survival

black cottonwood, growth/survival 38(4):27-30, Fall 1987

Damping-off

seed surface sterilization effects on 4 conifer species 38(3):18-21, Summer 1987

Deer Away 38(3):22-25, Summer 1987

Deer. See also Animal damage/control

repellants on Douglas-fir plantations 38(3):22-25, Summer 1987

Diseases. See also Fungi, fungal disease

brown spot in longleaf pine 38(3):16-17, Summer 1987

Fusarium cotyledon blight in ponderosa pine 38(3):18-21, Summer 1987

phytophthora root rot control 38(4):37-39, Fall 1987

Dormancy, bud

Douglas-fir 38(2):11-15, Spring 1987

interior spruce 38(2):11-15, Spring 1987

lodgepole pine 38(2):11-15, Spring 1987

Douglas-fir. See also Rocky Mountain Douglas-fir

cold storage of seedlings 38(2):11-15, Spring 1987

deer and elk, repellants on Douglas-fir plantations 38(3):22-25, Summer 1987

grafts

leader growth 38(4):14-18, Fall 1987

plagiotropism 38(4):14-18, Fall 1987

root stock age effects 38(4):14-18, Fall 1987

nutrient and metal concentrations in compostgrown seedlings 38(2):22-27, Spring 1987

plantation seedlings, effects of deer repellants

38(3):22-25, Summer 1987

seedling growth

response to *Pisolithus tinctorius*

inoculation 38(1):18-21, Winter 1987

sawdust-sludge compost 38(2):22-27, Spring 1987

seedling physiology, effect of cold storage 38(2):11-15, Spring 1987

seedlings, cold-stored

bud dormancy status 38(2):11-15, Spring 1987

carbohydrate reserves 38(2):11-15, Spring 1987

cold hardiness 38(2):11-15, Spring 1987

root growth potential 38(2):11-15, Spring 1987

water relations 38(2):11-15, Spring 1987

storage of seedlings

cold 38(2):11-15, Spring 1987

freezer 38(2):7-10, Spring 1987

Drought resistance. See Moisture stress

Eastern white pine

prechilling, extended, effect on seeds 38(4): 6-9, Fall 1987

seeds fungal infection 38(4):6-9, Fall 1987

germination 38(4): 6-9, Fall 1987

Economics

- cost control, nursery operations and genetically improved seed 38(3):3-8, Summer 1987
- feasibility of growing black cherry, black walnut, and northern red oak in northern California 38(4):31-36, Fall 1987

Elaeagnus umbellata. See Autumn olive

Elk. See also Animal damage/control

- repellants, Douglas-fir plantations 38(3):22-25, Summer 1987

Engelmann spruce seedling growth response to *Pisolithus tinctorius* inoculations 38(1):18-21, Winter 1987

Equipment

- mist blower, Hardi-Combi I 38(4):3-5, Fall 1987
- nursery, custom-built, list of 40 available construction drawings 38(2):3-6, Spring 1987
- scarification
 - Birch Quik-Tach Scalper 38(1):3-5, Winter 1987
 - cable-yarder 38(1):6-10, Winter 1987
 - Rocky Mountain scalper 38(1):3-5, Winter 1987
- Vexar plastic seedling protector 38(3):26-29, Summer 1987
- wire girdle, loblolly pine grafts 38(3):33-35, Summer 1987

Eucalyptus macarthurii

- aboveground biomass production 38(4):27-30, Fall 1987
- frost damage 38(4):27-30, Fall 1987
- seed source/provenance stock 38(4):27-30, Fall 1987
- survival and early growth 38(4):27-30, Fall 1987

Eucalyptus viminalis

- aboveground biomass production 38(4):27-30, Fall 1987
- frost damage 38(4):27-30, Fall 1987
- seed source/provenance stock 38(4):27-30, Fall 1987
- survival and early growth 38(4):27-30, Fall 1987

Fraser fir

- cold storage and seedling growth 38(2):19-21, Spring 1987
- container-seedling growth acceleration 38(2):19-21, Spring 1987
- seed source effect on seedling growth 38(2):19-21, Spring 1987

Freezer storage. See Seedling storage, cold/ frozen

Fumigation. See Soil treatment

Fungi, fungal disease. See also Fungicides

Fusarium species

- bleach-sterilized seeds of 4 conifer species 38(3):18-21, Summer 1987
- sycamore seedlings, cold-stored 38(1):26-30, Winter 1987
- infection in white spruce and eastern white pine seeds 38(4):6-9, Fall 1987
- microorganisms on cold-stored sycamore seedlings 38(1):26-30, Winter 1987
- Phytophthora* species, control of in Pacific Northwest nurseries 38(4):37-39, Fall 1987
- Pisolithus tinctorius*, inoculation effects on 5 Rocky Mountain conifer species 38(1):18-21, Winter 1987

Fungicides

- benomyl
 - loblolly pine seedlings 38(4):19-22, Fall 1987
 - sycamore seedlings 38(1):26-30, Winter 1987
- bleach, 40% laundry, 4 conifer species 38(3):18-21, Summer 1987
- metalaxyl vs. phytophthora root rot 38(4):37-39, Fall 1987
- propiconazol, sycamore seedlings 38(1):26-30, Winter 1987

Fusarium. See Fungi, fungal disease

Genetic improvement

- genetically improved seed for nursery operations 38(3):3-8, Summer 1987
- loblolly pine, joint efforts in southern U.S. 38(3):33-35, Summer 1987

Germination. See Seed germination

Grafting, Douglas-fir 38(4):14-18, Fall 1987

Greenhouse operations. See Nursery operations;
Equipment

Growing media. See *Soil* treatment

Growth. See Seedling growth/survival; Cuttings

Hand tools. See Equipment

Hardi-Combi I mist blower 38(4) :3-5, Fall 1987

Heavy metals. See Metals, heavy

History plots. See Nursery operations

IFA-PNW cooperative tree improvement program
38(3):3-8, Summer 1987

Implements. See Equipment

Insect damage/control

sawfly and black cherry 38(4):31-36, Fall 1987
termites, subterranean, and pecan seedlings
38(2):28-30, Spring 1987

Interior spruce

cold-stored seedlings and bud dormancy 38(2):11-15,
Spring 1987
cold hardiness 38(2):11-15, Spring 1987
root growth potential 38(2):11-15, Spring 1987

Juglans nigra. See Black walnut

Limber pine

seedling growth response to *Pisolithus tinctorius*
inoculation 38(1):18-21, Winter 1987

Liqua Gel

use on cold-stored sycamore seedlings
38(1):26-30, Winter 1987

Liriodendron tulipifera. See Yellow-poplar

Loblolly pine

droughty sites, outplanting on 38(4):19-22, Fall 1987
genetic improvement 38(3):33-35, Summer 1987
male flower production, effect of wire girdle on graft
38(3):33-35, Summer 1987
pollen distribution in seed orchard 38(4):3-5, Fall 1987

seed yield increase, effect of mist blower
38(4):3-5, Fall 1987

seedling survival-sowing date relationship
38(4):19-22, Fall 1987

seedlings, stored, survival affected by:
benomyl root dip 38(4):19-22, Fall 1987
seedling height 38(4):19-22, Fall 1987
shoot-to-root ratio 38(4):19-22, Fall 1987

Lodgepole pine

seedling growth response to *Pisolithus tinctorius*
inoculation 38(1):18-21, Winter 1987

seedlings, cold-stored bud dormancy 38(2):11-15, Spring
1987

cold hardiness 38(2):11-15, Spring 1987

freezer storage practices 38(2):7-10, Spring 1987

root growth potential 38(2):11-15, Spring 1987

Longleaf pine

brown spot 38(3):16-17, Summer 1987

seedling size-survival/growth 38(3):16-17, Summer 1987

Lorsban 4E

vs. subterranean termites in pecan seedlings
38(2):28-30, Spring 1987

Lycopersicon esculentum* var. *pelican. See
Tomato

Machinery. See Equipment

Metals, heavy

concentrations in roots of sawdust-sludge-composted
conifer seedlings 38(2):22-27, Spring 1987

Mine spoil reclamation

autumn olive 38(1):14-17, Winter 1987

Moisture retention

sycamore, Liqua-Gel 38(1):26-30, Winter 1987

tomato seedlings, Terra-sorb 38(1):24-25, Winter 1987

Moisture stress

antitranspirants, 3 film-forming, effect
on 38(4):23-26, Fall 1987

- black spruce outplanted seedlings 38(4):23-26, Fall 1987
 cold-stored western conifer seedlings 38(2):11-15, Spring 1987
 loblolly pine 38(4):19-22, Fall 1987
 northern California—black walnut, black cherry, northern red oak 38(4):31-36, Fall 1987
- Mycorrhizae**
 ectomycorrhizal association with stunting of white spruce 38(1):22-23, Winter 1987
 Pisolithus tinctorius, inoculation of 5 Rocky Mountain conifer species 38(1):18-21, Winter 1987
- Myocaster coypu.** See Nutria
- Noble fir**
 freezer-storage practices 38(2):7-10, Spring 1987
 growth in sawdust-sludge compost 38(2):22-27, Spring 1987
 nutrient/heavy metal concentrations 38(2):22-27, Spring 1987
- North Carolina State University-Industry Tree Improvement Cooperative** 38(3):33-35, Summer 1987
- Northern red oak**
 adaptability to northern California and 10-year survival/growth results 38(4):31-36, Fall 1987
 pocket gopher/soil moisture major factors 38(4):31-36, Fall 1987
- Nursery operations.** See also Equipment
 antitranspirants, black spruce seedlings 38(4):23-26, Fall 1987
 chilling, Fraser fir seedlings 38(2):19-21, Spring 1987
 cold hardiness, simple method for evaluating whole plant 38(2):16-18, Spring 1987
 cold storage
 Douglas-fir seedlings 38(2):11-15, Spring 1987
 lodgepole pine seedlings 38(2):11-15, Spring 1987
 interior spruce seedlings 38(2):11-15, Spring 1987
 compost, sawdust-sludge, for 3 conifer species 38(2):22-27, Spring 1987
 freezer-storage practices for western conifer seedlings 38(2):7-10, Spring 1987
 fungal disease control
 eastern white pine 38(4):6-9, Fall 1987
 phytophthora root rot 38(4):37-39, Fall 1987
 sycamore 38(1):26-30, Winter 1987
 western conifers 38(3):18-21, Summer 1987
 white spruce 38(4):6-9, Fall 1987
 germination recommendations
 Austrian pine 38(3):18-21, Summer 1987
 autumn olive 38(1):14-17, Winter 1987
 eastern white pine 38(4):6-9, Fall 1987
 pin cherry 38(3):30-32, Summer 1987
 ponderosa pine 38(3):18-21, Summer 1987
 Rocky Mountain Douglas-fir 38(3):18-21, Summer 1987
 Scotch pine 38(3):18-21, Summer 1987
 white spruce 38(4):6-9, Fall 1987
 history plots
 applications to nursery management 38(3):9-15, Summer 1987
 installation 38(3):9-15, Summer 1987
 monitoring 38(3):9-15, Summer 1987
 seedling growth curves and phenology 38(3):9-15, Summer 1987
 sowing rate factors 38(3):9-15, Summer 1987
 longitudinal splitting, propagation technique, yellow poplar 38(4):10-13, Fall 1987
 management 38(3):9-15, Summer 1987
 packaging, freezer-stored western conifers 38(2):7-10, Spring 1987
 phytophthora root rot management in Pacific Northwest conifer nurseries 38(4):37-39, Fall 1987

Pisolithus tinctorius inoculation of conifer container seedlings 38(1):18-21, Winter 1987
prechilling, extended, of white spruce and eastern white pine
fungal infection of seeds 38(4):6-9, Fall 1987
germination 38(4):6-9, Fall 1987
seed, surface sterilization with bleach 38(3):18-21, Summer 1987
seed-use efficiency improvement 38(3):9-15, Summer 1987
soil treatment
crusting prevention, ponderosa pine 38(1):11-13, Winter 1987
fumigation, white spruce 38(1):22-23, Winter 1987
water-holding capacity soil amendment 38(1):24-25, Winter 1987
sowing 38(3):9-15, Summer 1987
storage
cold, of conifer seedlings 38(2):11-15, Spring 1987
freezer, practices for western conifer seedlings 38(2):7-10, Spring 1987
termite prevention, pecan seedlings 38(2):28-30, Spring 1987
tree improvement
implications for nursery management 38(3):3-8, Summer 1987
key leverage points for nursery system 38(3):3-8, Summer 1987
seed, genetically improved 38(3):3-8, Summer 1987
seedling quality improvement 38(3):9-15, Summer 1987

Nutria. See also Animal damage/control
baldcypress seedling damage 38(3):26-29, Summer 1987
control trial with Vexar seedling protectors 38(3):26-29, Summer 1987

Nutrient concentration levels

carbohydrate reserves, cold-stored Douglas fir 38(2):11-15, Spring 1987

in 3 compost-grown conifer species 38(2):22-27, Spring 1987

Pacific Northwest

Eucalyptus macarthurii and *E. viminalis* 38(4):27-30, Fall 1987

Packaging practices

freezer-stored western conifers 38(2):7-10, Spring 1987

Pecan termites, subterranean, damage to seedlings 38(2):28-30, Spring 1987

Pest control. See Animal damage/control; Insect damage/control

Phytophthora root rot

chemical management—fumigants, fungicides, chlorination, metalaxyl 38(4):37-39, Fall 1987
cultural management—water, sanitation, resistant species 38(4):37-39, Fall 1987

Picea

engelmannii. See Engelmann spruce
glauca. See White spruce
glauca-engelmannii complex. See Interior spruce
mariana. See Black spruce

Pin cherry

seed germination and temperature fluctuation influence 38(3):30-32, Summer 1987

Pinus

contorta. See Lodgepole pine
flexilis. See Limber pine
nigra. See Austrian pine
palustris. See Longleaf pine
ponderosa. See Ponderosa pine
strobus. See Eastern white pine
sylvestris. See Scotch pine
taeda. See Loblolly pine

Pisolithus tinctorius

Douglas-fir 38(1):18-21, Winter 1987
Engelmann spruce 38(1):18-21, Winter 1987
limber pine 38(1):18-21, Winter 1987

lodgepole pine 38(1):18-21, Winter 1987

ponderosa pine 38(1):18-21, Winter 1987

Plagiotropism

Douglas-fir grafts 38(4):14-18, Fall 1987

Plant moisture stress. See also Moisture stress

outplanted black spruce seedlings 38(4):23-26, Fall 1987

Platanus occidentalis. See Sycamore

Pocket gopher damage. See also Animal damage/ control

black cherry in northern California 38(4):31-36, Fall 1987

northern red oak in northern California 38(4):31-36, Fall 1987

Pollen

distribution

guidelines for use of mist blower 38(4):3-5, Fall 1987

loblolly pine, effects of mist blower 38(4):3-5, Fall 1987

production

loblolly pine, effects of wire girdle 38(3):33-35, Summer 1987

Ponderosa pine

Fusarium and damping-off 38(3):18-21, Summer 1987

germination of bleach surface-sterilized seed

38(3):18-21, Summer 1987

heavy metal concentrations in compost-grown seedlings

38(2):22-27, Spring 1987

nutrient concentrations in compost-grown seedlings

38(2):22-27, Spring 1987

seedling freezer-storage practices 38(2):7-10, Spring

1987

seedling growth

Pisolithus tinctorius inoculations, response to 38(1):18-21, Winter 1987

sawdust-sludge compost 38(2):22-27, Spring 1987

seed covering materials, test results 38(1):11-13, Winter 1987

Populus trichocarpa. See Black cottonwood

Propiconazol

use on cold-stored sycamore seedlings

38(1):26-30, Winter 1987

Provenance. See Seed source

Prunus

pennsylvanica. See Pin cherry

serotina. See Black cherry

Pseudotsuga menziesii. See Douglas-fir

Pseudotsuga menziesii var. glauca. See Rocky

Mountain Douglas-fir

Quercus rubra. See Northern red oak

Red alder

biomass production, aboveground 38(4):27-30, Fall 1987

early height growth 38(4):27-30, Fall 1987

frost tolerance 38(4):27-30, Fall 1987

seedling survival 38(4):27-30, Fall 1987

Reforestation

Pacific Northwest tree improvement status

38(3):3-8, Summer 1987

seed orchards role 38(3):3-8, Summer

1987; 38(3):33-35, Summer 1987

site preparation equipment 38(1):3-5, Winter

1987; 38(1):6-10, Winter 1987

Repelliff

use on deer and elk 38(3):22-25, Summer 1987

Reticulitermes flavipes. See Termites,

subterranean

RGP. See Root growth potential

Rocky Mountain Douglas-fir. See also Douglas-fir

Fusarium and damping-off 38(3):18-21, Summer 1987

germination of bleach surface-sterilized

seed 38(3):18-21, Summer 1987

Rocky Mountain Scalper 38(1):3-5, Winter 1987

Root collar diameter

effect on longleaf pine seedling

growth 38(3):16-17, Summer 1987

Root disease

Fusarium species 38(3):18-21, Summer 1987
phytophthora root rot 38(4):37-39, Fall 1987

Root growth potential, cold-stored seedlings

Douglas-fir 38(2):11-15, Spring 1987
interior spruce 38(2):11-15, Spring 1987
lodgepole pine 38(2):11-15, Spring 1987

Rootstock age

effect on Douglas-fir grafts 38(4):14-18, Fall 1987

Scarification equipment 38(1):3-5, Winter 1987; 38(1):6-10, Winter 1987

Scirrhia acicola. See Brown spot

Scotch pine

Fusarium and damping-off 38(3):18-21, Summer 1987
germination of bleach surface-sterilized seed 38(3):18-21, Summer 1987

Seed, genetically improved

nursery manager, implications for 38(3):3-8, Summer 1987
Pacific Northwest, review of 38(3):3-8, Summer 1987
production system economics 38(3):3-8, Summer 1987

Seed germination

Austrian pine, bleach sterilization treatment 38(3):18-21, Summer 1987
autumn olive, stratification/temperature recommendations 3(1):14-17, Winter 1987
eastern white pine, effect of prechilling 38(4):6-9, Fall 1987
pin cherry, recommendations for 38(3):30-32, Summer 1987
ponderosa pine, bleach sterilization treatment 38(3):18-21, Summer 1987
Rocky Mountain Douglas-fir, bleach sterilization treatment 38(3):18-21, Summer 1987
Scotch pine, bleach sterilization treatment 38(3):18-21, Summer 1987
white spruce, effect of prechilling 38(4):6-9, Fall 1987

Seed orchard

Douglas-fir region 38(3):3-8, Summer 1987
loblolly pine 38(3):33-35, Summer 1987; 38(4):3-5, Fall 1987

Seed source

Eucalyptus 38(4):27-30, Fall 1987
Fraser fir 38(2):19-21, Spring 1987

Seed, sowing

family 38(3):3-8, Summer 1987
loblolly pine, time of 38(4):19-22, Fall 1987
recommendations based on history plots 38(3):9-15, Summer 1987

Seed treatment. See also Seed germination

coverings, effect on ponderosa pine 38(1):11-13, Winter 1987
prechilling, extended
eastern white pine 38(4):6-9, Fall 1987
germination/fungal infection, effects on 38(4):6-9, Fall 1987
white spruce 38(4):6-9, Fall 1987
sterilization, surface, with bleach, of 4 conifer species 38(3):18-21, Summer 1987
stratification
autumn olive 38(1):14-17, Winter 1987
pin cherry 38(3):30-32, Summer 1987
temperature
autumn olive 38(1):14-17, Winter 1987
pin cherry 38(3):30-32, Summer 1987

Seedling growth/survival

baldcypress 38(3):26-29, Summer 1987
black cherry 38(4):31-36, Fall 1987
black spruce 38(4):23-26, Fall 1987
black walnut 38(4):31-36, Fall 1987
Douglas-fir 38(1):18-21, Winter 1987;
38(2):7-10, Spring 1987; 38(2):22-27, Spring 1987; 38(3):22-25, Summer 1987
Engelmann spruce 38(1):18-21, Winter 1987
Eucalyptus macarthurii 38(4):27-30, Fall 1987
Eucalyptus viminalis 38(4):27-30, Fall 1987
Fraser fir 38(2):19-21, Spring 1987
limber pine 38(1):18-21, Winter 1987
loblolly pine 38(4):19-22, Fall 1987
lodgepole pine 38(1):18-21, Winter

1987; 38(2):7-10, Spring 1987
 longleaf pine 38(3):16-17, Summer 1987
 noble fir 38(2):7-10, Spring 1987; 38(2):22-27, Spring 1987
 northern red oak 38(4):31-36, Fall 1987
 pecan 38(2):28-30, Spring 1987
 ponderosa pine 38(1):11-13, Winter 1987; 38(1):18-21, Winter 1987; 38(2):7-10, Spring 1987; 38(2):22-27, Spring 1987
 red alder 38(4):27-30, Fall 1987
 sycamore 38(1):26-30, Winter 1987
 tomato 38(1):24-25, Winter 1987
 western hemlock 38(2):7-10, Spring 1987
 white spruce 38(1):22-23, Winter 1987
 yellow poplar 38(4):10-13, Fall 1987

Seedling storage, cold/frozen

method

Fraser fir 38(2):19-21, Spring 1987
 loblolly pine 38(4):19-22, Fall 1987
 sycamore 38(1):26-30, Winter 1987
 western conifers 38(2):7-10, Spring 1987

seedling physiology, effects on

Douglas-fir 38(2):11-15, Spring 1987
 interior spruce 38(2):11-15, Spring 1987
 lodgepole pine 38(2):11-15, Spring 1987

Site preparation

scarification equipment 38(1):3-5, Winter 1987; 38(1):6-10, Winter 1987
 steep slopes, cable-yarder alternative 38(1):610, Winter 1987

Soil moisture. See Moisture stress**Soil treatment**

amendment, tomato water-holding capacity 38(1):24-25, Winter 1987
 crusting prevention in nurseries, ponderosa pine 38(1):11-13, Winter 1987
 fumigation-mycorrhizae-stunting relationship, white spruce 38(1):22-23, Winter 1987
 sawdust-sludge compost, 3 conifer species 38(2):22-27, Spring 1987

Sowing. See Seed, sowing**Storage.** See Seedling storage, cold/frozen

Stratification. See Seed treatment; Seed germination

Stunting

white spruce 38(1):22-23, Winter 1987

Survival. See Seedling growth/survival; Cuttings

Sycamore

cold-stored seedlings and chemical treatment effects 38(1):26-30, Winter 1987
 dieback 38(1):26-30, Winter 1987

Taxodium distichum. See Baldcypress

Termites

subterranean, damage to pecan seedlings 38(2):28-30, Spring 1987

Terra-sorb

use in container tomatoes 38(1):24-25, Winter 1987

Thomomys spp. See Pocket gopher

Tomato

container seedlings, water-holding capacity 38(1):24-25, Winter 1987
 wilting, effect of Terra-sorb 38(1):24-25, Winter 1987

Tools. See Equipment

Tree improvement

Pacific Northwest, review and implications for nursery manager 38(3):3-8, Summer 1987
 South, loblolly pine genetic base 38(3):33-35, Summer 1987

Tsuga heterophylla. See Western hemlock

Vapor Gard

effect on cold-stored sycamore seedlings 38(1):26-30, Winter 1987

Vexar seedling protector

effect on baldcypress 38(3):26-29, Summer 1987

Water. See *also* Moisture retention; Moisture stress management practices vs. phytophthora root rot 38(4):37-39, Fall 1987
 relationship with cold-stored conifer seedlings 38(2):11-15, Spring 1987

testing for contamination by *Phytophthora*
species 38(4):37-39, Fall 1987

Western hemlock

freezer-storage practices for seedlings 38(2):7-10,
Spring 1987

White spruce

mycorrhizal fungi and soil fumigation 38(1):22-23, Winter
1987

seeds

fungal infection 38(4):6-9, Fall 1987

germination 38(4):6-9, Fall 1987

prechilling, extended 38(4):6-9, Fall 1987

stunting, ectomycorrhizal deficiency 38(1):22-23, Winter
1987

Wilt-pruf NCF

black spruce seedlings 38(4):23-26, Fall 1987

X2-1337

black spruce seedlings 38(4):23-26, Fall 1987

Yellow-poplar

propagation technique 38(4):10-13, Fall 1987

seedlings, longitudinally split, survival 38(4):10-
13, Fall 1987