Application Of Somatic Embryogenesis In The Production Of Christmas Trees

Ulrika Egertsdotter¹, Harald Kvaalen², Ola Gram Daehlen³ and Maigull Appelgren⁴

¹Institute of Paper Science and technology, 500 10th Street, N.W. Atlanta, GA 30318 ²Norwegian Forest Research Institute, Hogskoleveien 12, N-1432 Aas, Norway ³Oppland Skogselskap, Biri Skogplanteskole, N-2820 Biri, Norway ⁴Agricultural University of Norway, Department of Horticulture and Crop Sciences, N-1432 Aas, Norway

Ulrika.Egertsdotter@ipst.edu

A growing area of interest within the forestry industry is the production of Christmas trees and decoration greenery. Abies lasiocarpa is one of the Abies-species which is suitable for Christmas tree production due to their long-lasting greenery. In Norway, the trees in the populations of A. lasiocarpa show a great diversity in morphology and growth. The lack of uniformity between individuals has thus prompted for the use of efficient clonal propagation methods for enabling the exploitation of valuable genotypes in plantation forestry of this species. In vitro propagation by somatic embryogenesis has been successfully achieved for many coniferous species. The use of somatic embryogenesis for plant propagation of A. lasiocarpa was evaluated. Protocols, and in particular the culture media, for somatic embryogenesis in conifers has predominantly been developed for *Picea*-species and today plantlets can readily be produced from most species. The obstacles earlier encountered with the Abies-species, indicate differences in embryo physiology in Abies-and Picea-species. We have designed a novel growth medium for initiation of somatic embryogenesis based on the natural environment of the zygotic embryo. The interactions between the media composition, the developmental stage of the explants, and the influence of the genetic background were tested. We have also explored the positive effects of microbial compounds on growth and development of plantlets derived from somatic embryos.